

Human Immunodeficiency Virus (HIV)

Prognosis Including Monitoring

CPT: 87536, 87539

Medically Supportive ICD Codes are listed
on subsequent page(s) of this document

CMS National Coverage Policy

Coverage Indications, Limitations, and/or Medical Necessity

HIV quantification is achieved through the use of a number of different assays which measure the amount of circulating viral RNA. Assays vary both in methods used to detect viral RNA as well as in ability to detect viral levels at lower limits. However, all employ some type of nucleic acid amplification technique to enhance sensitivity, and results are expressed as the HIV copy number.

Quantification assays of HIV plasma RNA are used prognostically to assess relative risk for disease progression and predict time to death, as well as to assess efficacy of anti-retroviral therapies over time.

HIV quantification is often performed together with CD4+ T cell counts which provide information on extent of HIV induced immune system damage already incurred.

Indications

1. A plasma HIV RNA baseline level may be medically necessary in any patient with confirmed HIV infection.
2. Regular periodic measurement of plasma HIV RNA levels may be medically necessary to determine risk for disease progression in an HIV-infected individual and to determine when to initiate anti-retroviral treatment regimens.
3. In clinical situations where risk of HIV infection is significant and initiation of therapy is anticipated, a baseline HIV quantification may be performed. These situations include:
 - a) Persistence of borderline or equivocal serologic reactivity in an at-risk individual.
 - b) Signs and symptoms of acute retroviral syndrome characterized by fever, malaise, lymphadenopathy and rash in an at-risk individual.

Limitations

1. Viral quantification may be appropriate for prognostic use including baseline determination, periodic monitoring, and monitoring of response to therapy. Use as a diagnostic test method is not indicated.
2. Measurement of plasma HIV RNA levels should be performed at the time of establishment of an HIV infection diagnosis. For an accurate baseline, 2 specimens in a 2-week period are appropriate.
3. For prognosis including anti-retroviral therapy monitoring, regular, periodic measurements are appropriate. The frequency of viral load testing should be consistent with the most current Centers for Disease Control and Prevention guidelines for use of anti-retroviral agents in adults and adolescents or pediatrics.
4. Because differences in absolute HIV copy number are known to occur using different assays, plasma HIV RNA levels should be measured by the same analytical method. A change in assay method may necessitate re-establishment of a baseline.
5. Nucleic acid quantification techniques are representative of rapidly emerging & evolving new technologies. Users advised to remain current on FDA-approval status.

Visit [SonoraQuest.com/Medicare](https://www.SonoraQuest.com/Medicare) to view current limited coverage tests, reference guides, and policy information.

To view the complete policy and the full list of medically supportive codes, please refer to the CMS website reference

Human Immunodeficiency Virus (HIV)

Prognosis Including Monitoring

CPT: 87536, 87539

The ICD10 codes listed below are the top diagnosis codes currently utilized by ordering physicians for the limited coverage test highlighted above that are also listed as medically supportive under Medicare's limited coverage policy. **If you are ordering this test for diagnostic reasons that are not covered under Medicare policy, an Advance Beneficiary Notice form is required.**

***Note—Boded diagnoses below have the highest utilization**

Please refer to the Limitations or Utilization Guidelines section on previous page(s) for frequency information.

Code	Description
B20	Human immunodeficiency virus [HIV] disease
B97.35	Human immunodeficiency virus, type 2 [HIV 2] as the cause of diseases classified elsewhere
R75	Inconclusive laboratory evidence of human immunodeficiency virus [HIV]
Z21	Asymptomatic human immunodeficiency virus [HIV] infection status

Visit [SonoraQuest.com/Medicare](https://www.cms.gov/Medicare) to view current limited coverage tests, reference guides, and policy information.

To view the complete policy and the full list of medically supportive codes, please refer to the CMS website reference

www.cms.gov ►

Last updated: 10/01/21

Disclaimer:

This diagnosis code reference guide is provided as an aid to physicians and office staff in determining when an ABN (Advance Beneficiary Notice) is necessary. Diagnosis codes must be applicable to the patient's symptoms or conditions and must be consistent with documentation in the patient's medical record. Sonora Quest Laboratories does not recommend any diagnosis codes and will only submit diagnosis information provided to us by the ordering physician or his/her designated staff. The CPT codes provided are based on AMA guidelines and are for informational purposes only. CPT coding is the sole responsibility of the billing party. Please direct any questions regarding coding to the payer being billed.

SonoraQuest.com

Sonora Quest Laboratories, any associated logos, and all associated Sonora Quest Laboratories registered or unregistered trademarks are the property of Sonora Quest Laboratories. All third-party marks—® and ™—are the property of their respective owners. © 2021 Sonora Quest Laboratories. All rights reserved.