



OmegaCheck

CPT Code **82542**

Test Code **906900**

Sample Type **Whole Blood**

Tube Type **Lavender-Top**

Low omega-3 fatty acid levels are associated with:

- Hypertriglyceridemia
- High blood pressure
- Increased risk of heart disease

Description

Omega-3 and omega-6 fatty acids are polyunsaturated long chain fatty acids (PUFA) required by the body for proper functioning, normal growth and the formation of neural synapses and cellular membranes. Omega-3 and -6 fatty acids are considered “essential” and obtained primarily from dietary sources.

Three of the most important omega-3 fatty acids are eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA) and docosahexaenoic acid (DHA). Omega-3 fatty acids are primarily obtained from food sources, such as oily fish. They have antioxidant¹, anti-inflammatory² and anti-thrombotic³ effects, and can help to reduce triglyceride levels⁴⁻⁶. Two of the most important omega-6 fatty acids are arachidonic acid (AA) and linoleic acid (LA). Omega-6 fatty acids are obtained from animal sources and plant oils, and have pro-inflammatory^{2,7} and pro-thrombotic⁷ properties at high levels.

Clinical Use

OmegaCheck™ may be performed on individuals with hypercholesterolemia, hypertriglyceridemia, hypertension, and/or those at high metabolic or cardiovascular risk.

Clinical Significance

- Consumption of omega-3 fatty acids reduces the occurrence of major acute cardiac events in healthy individuals or patients with cardiovascular risk factors or who have cardiovascular disease⁸⁻¹⁴.
- Consumption of omega-3 fatty acids leads to a reduction in triglycerides⁴⁻⁶ and non-HDL⁶, as well as Lp-PLA₂ levels⁶.
- A high intake of omega-6 fatty acid precursors can interfere with the absorption of omega-3 fatty acids⁸.
- The mean omega-6:omega-3 ratio of the standard American diet is approximately 10:1⁸. A diet with an omega-6:omega-3 fatty acid ratio of 4:1 or less may reduce total mortality up to 70% over 2 years¹¹.

Testing Frequency

Testing frequency depends on the individual's medical history. OmegaCheck™ may be run alongside a standard lipid panel or other cardiometabolic tests.

Sample Type

OmegaCheck™ should be performed on a whole blood sample. Fasting samples are preferred, but not required, and omega-3 supplementation should not be altered immediately prior to the blood draw.

Commercial Insurance or Medicare Coverage

Coverage guidelines, also known as NCD (National Coverage Determination) or LCD (Local Coverage Determination), have not been established or posted by CMS (Medicare & Medicaid). We have reviewed the larger Carriers (Aetna, UnitedHealthcare, Cigna, Blues) and information has not been posted or is limited. Medical necessity and specificity of diagnosis should be provided when ordering this test.

Understanding Medical Necessity

The following ICD-10 codes for OmegaCheck™ are listed as a convenience for the ordering physician. The ordering physician should report the diagnosis code that best describes the reason for performing the test.

Diagnosis	Diagnosis Code
Type 2 Diabetes Mellitus with Hyperglycemia	E11.65
Type 2 Diabetes Mellitus without Complications	E11.9
Other Specified Diabetes Mellitus without Complications	E13.9
Pure Hypercholesterolemia, Unspecified	E78.00
Familial Hypercholesterolemia	E78.01
Pure Hyperlipidemia	E78.1
Mixed Hyperlipidemia	E78.2
Other Hyperlipidemia	E78.4
Hyperlipidemia, Unspecified	E78.5
Metabolic Syndrome	E88.81
Essential (primary) Hypertension	I10
Atherosclerotic Heart Disease of Native Coronary Artery without Angina Pectoris	I25.10
Impaired Fasting Glucose	R73.01
Impaired Glucose Tolerance Test (oral)	R73.02

References

1. Kesavulu MM et al. Effect of ω -3 fatty acids on lipid peroxidation and antioxidant enzyme status in type 2 diabetic patients. *Diabetes Metab.* 2002; 28: 20-26.
2. James MJ et al. Dietary polyunsaturated fatty acids and inflammatory mediator production. *Am J Clin Nutr.* 2000; 71: 343s-348s.
3. Engstrom K et al. Effect of low-dose aspirin in combination with stable fish oil on whole blood production of eicosanoids. *Prostaglandins Leukot Essent Fatty Acids.* 2001; 64: 291-297.
4. Balk E et al. Effects of omega-3 fatty acids on cardiovascular risk factors and intermediate markers of cardiovascular disease. *Evid Rep Technol Assess* 2004; Mar(93): 1-6.
5. Musa-Veloso K et al. Long-chain omega-3 fatty acids eicosapentaenoic acid and docosahexaenoic acid dose-dependently reduce fasting serum triglycerides. *Nutrition Reviews.* 2010; 68: 155-167.
6. Kastlein JJP et al. Omega-3 free fatty acids for the treatment of severe hypertriglyceridemia: The EpanoVa for Lowering Very high triglyceridEs (EVOLVE) trial. *J Clin Lipidol.* 2014; 8: 94-106.
7. Schmitz G. The opposing effects of n-3 and n-6 fatty acids. *Prog Lipid Res.* 2008; 47: 147-155.
8. Saito Y et al. Effects of EPA on coronary artery disease in hypercholesterolemic patients with multiple risk factors: Sub-analysis of primary prevention cases from Japan EPA Lipid Intervention Study (JELIS). *Atherosclerosis.* 2008; 200: 135-140.
9. Marchioli R et al. Early protection against sudden death by n-3 polyunsaturated fatty acids after myocardial infarction. Time-course analysis of the results of the Gruppo-Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico (GISSI)-Prevenzione. *Circulation.* 2002; 105: 1897-1903.
10. Pottala JV et al. Blood eicosapentaenoic and docosahexaenoic acids predict all-cause mortality in patients with stable coronary heart disease: The Heart and Soul Study. *Circ Cardiovasc Qual Outcomes.* 2010; 3: 406-412.
11. de Lorgeril M et al. Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet.* 1994; 343: 1454-1459.
12. Simopoulos AP. The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Exp Biol Med.* 2009; 233: 674-688.
13. Albert CM et al. Blood levels of long-chain n-3 fatty acids and the risk of sudden death. *N Engl J Med.* 2002; 346: 1113-1118.
14. Kesavulu MM et al. Effect of ω -3 fatty acids on lipid peroxidation and antioxidant enzyme status in type 2 diabetic patients. *Diabetes Metab.* 2002; 28: 20-26.

SonoraQuest.com

Sonora Quest Laboratories, any associated logos, and all associated Sonora Quest Laboratories registered or unregistered trademarks are the property of Sonora Quest Laboratories. All third-party marks —© and ™—are the property of their respective owners. © 2019 Sonora Quest Laboratories. All rights reserved. 4/2019

6701 Carnegie Ave. | Suite 500 | Cleveland, OH 44103
p 866.358.9828 | f 866.869.0148
clevelandheartlab.com | knowyourrisk.com | chlcm.com

RELATIVE RISK

OmegaCheck™
(% by weight)

≥5.5
Low

3.8-5.4
Moderate

≤3.7
High

The OmegaCheck™ was developed and validated at Cleveland HeartLab with the support of Nutrasource Diagnostics, Inc.

Sample Considerations

Omega-3 and -6 fatty acid levels can be measured in whole blood or within red blood cell (RBC) membranes. The OmegaCheck™ test measures omega-3 and -6 fatty acid levels in whole blood. The whole blood test provides a complete picture of the amount of omega-3 and -6 fatty acids in the body, and may reflect more recent levels of supplementation or dietary intake. RBC membrane levels provide a picture of consumption levels over a longer period of time, because the incorporation of these fatty acids into the membranes of RBC takes several days to weeks. Regardless, both whole blood (EPA+DPA+DHA (%)) and RBC membrane (EPA + DHA (%)) measurements demonstrate positive correlation ($r=0.91$, $p<0.0001$)¹⁴. This finding has been confirmed by studies run by Cleveland HeartLab.

Treatment Considerations

These treatment considerations are for educational purposes only. Specific treatment plans should be provided and reviewed by the treating practitioner.

✓ Assess dietary intake of omega-3 and omega-6 fatty acids.

- Dietary sources of omega-3 fatty acids include fatty fishes, such as salmon or sardines, nuts, and plant oils. Foods high in omega-6 fatty acids include red meat, poultry, eggs, plant oils, and nuts.

✓ Consider omega-3 fatty acid supplementation.

- If currently taking, consider adjusting dosage and retest in 1-2 months.

✓ Assess lifestyle habits.

- Consider diet/exercise/weight reduction efforts if appropriate.